























Recommended methods for the Microbiological Examination of Food APHA, Inc. New York. Pp. 53-59.

[3] AOAC 2002. Official Method of analysis 12th Ed. Method 2.049 and 14.006. Association of Official Analytical Chem., Washington, DC. Pp. 450-451, 520-551 USA.

[4] Banureka V.D. and Mahendran T., 2009. Formulation of Whea-Soybean Biscuits And Their Quality Characteristics. Tropical Agricultural Research & Extension, 12(2): 62-66.

[5] Gifford, S.R., & Clydesdale, F.M. 1986. The psychophysical relationship between color and sodium chloride concentrations in model systems. Journal of Food Protection, 49(12), 977-982.

[6] Manley, D. 2000 Technology of biscuits, crackers and cookies. Woodhead Publishing, Cambridge.

[7] Oluwamukomi M.O., Oluwalana I.B., and Akinbowale O.F., 2011. Physicochemical and sensory properties of wheat cassava composite biscuit enriched with soy flour. African Journal of

Food Science , 5 (2) : 50 - 56.

[8] Operation Manual. Glutomatic System. Parten Instruments. ICC Standard No. 155 and No 158. AACC Method 38-12.

[9] Patki, P. E. and S. S. Arya. 1994. Studies on the development and storage stability of instant dhals. Indian Food Packer. 48:31-39.

[10] Rangana, S. 1991. Hand book of analysis and quality control for fruit and vegetable products, 2nd Ed. Pun. Tata McGrawHill Publishing Company Ltd., New Delhi.

[11] Ranhotra GS (1980). Nutritional profile of high protein cookies. Cereal Foods World 25 308-309.

[12] Rockland, L. B. and S. K. Nishi. 1980. Influence of water activity on food product quality and stability. Food Technol. 34:42-59.

[13] Ugwuona F.U., 2009. Chemical And Sensory Evaluation Of Soy-Fortified Cassava wheat Biscuit. Journal of Tropical Agriculture, Food, Environment and Extension, 8 (1): 55-59.

IJSER