















- Comparative effects of osmotic and ionic stresses. *Plant Prod. Sci.*, 10: 159-170.
- [9] Ashraf, M. and P.J.C. Harris. 2004. Potential biochemical indicators of salinity tolerance in plants. *Plant Sci.*, 166: 3-16
- [10] Zang, X. and S. Komatsu. 2007. A proteomics approach for identifying osmotic-stress-related proteins in rice. *Phytochemical*. 68: 426-437.
- [11] Karp, A., H. Steel, S. Parmar, K. Jones, R. Shewry and A. Breiman, 1987. Relative stability among barley plants regenerated from cultured immature embryos. *Genome*, 29: 405-412.
- [12] Karp, A., 1995. Somaclonal variation as a tool for crop improvement. *Euphytica*, 85: 295-302
- [13] Ganeshan, S., Baga, M., Harwey, B.L., Rosnagel, B.G., Scoles, G.J., Chibbar, R.N. (2003).
- [14] Seraj, Z.I., Islam, Z., Faruque, M.O., Devi, T., Ahmed, S. (1997). Identification of the regeneration potential of embryo derived calli from various Indica rice varieties. *Plant Cell Tissue and Organ Culture* 48, 9-13
- [15] Mahajan, S. and N. Tuteja. 2005. Cold, salinity and drought stresses: An overview. *Arch. Biochem. Biophys.*, 444: 139-158. Varieties differing in salinity resistance. *Plant Growth Regulation* 19, 2-218.
- [16] Khalequzzaman, M., Haq, N., Hoque, ME., Aditya, TL. (2005). Regeneration efficiency and genotypic effect of 15 Indica type Bangladeshi rice (*Oryza sativa* L.) landraces. *Plant Tissue Culture* 15, 33-42
- [17] Al-Bahrany, A.M. (2002). Callus growth and proline accumulation in response to Polyethylene glycol induced osmotic stress in rice *Oryza sativa* L. *Pakistan Journal of Biological Sciences* 15, 1294-1296.
- [18] O'Toole, J.C. (1982). Adaptation of rice to drought prone environments. In: *Drought resistant crops, with emphasis on rice*, IRRI, Los Baños.
- [19] Rashid, H., Yokoi, S., Toriyama, K., Hinata, K. (1996). Transgenic plant production mediated by *Agrobacterium* in Indica rice. *Plant Cell Reports* 15, 727-730.
- [20] Wang, W., B. Vinocur and A. Altman. 2003. Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta*, 218: 1-14.
- [21] Wang, W., B. Vinocur, O. Shoseyov and A. Altman. 2001. Biotechnology of plant osmotic stress tolerance: physiological and molecular considerations. *Acta Hort.*, 560: 285-292.
- [22] Wani, S.H., Gosal, S.S. (2011). Introduction of *OsglyII* gene into Indica rice through particle bombardment for increased salinity tolerance. *Biologia Plantarum* (in press).
- [23] Lutts, S., M. Almansouri and J.M. Kinet. 2004. Salinity and water stress have contrasting effects on the relationship between growth and cell viability during and after stress exposure in durum wheat callus. *Plant Sci.*, 167: 9-18.
- [24] Zalc, J.M., Wicr, H.B., Kidwell, K.K., Steber, CM. (2004). Callus induction and plant regeneration from mature embryos of a diverse set of wheat genotypes. *Plant Cell Tissue and Organ Culture* 76, 277-281.